

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/365184182>

The influence of the vacuum drying method on the physics chemical properties of stingless bee honey

Article · October 2022

CITATIONS

0

READS

65

4 authors, including:



Imron Meechai

Yala Rajabhat University

8 PUBLICATIONS 65 CITATIONS

SEE PROFILE



Isma-Ae Chelong

Yala Rajabhat University

13 PUBLICATIONS 43 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Utilization of Natural Product from stingless bee and edible and medicinal plant [View project](#)



CO-INFECTION WITH COXIELLA-LIKE BACTERIA AND BABESIA IN GOAT TICKS FROM SOUTHERN THAILAND [View project](#)

The Influence of the Vacuum drying method on the physicochemical properties of Stingless bee Honey

I. Meechai^{1*}, R. Yuso², I. Chelong³, A. Sukswan⁴

^{1,2}Chemistry Program, Faculty of Science Technology and Agriculture, Yala Rajabhat University, Yala, Thailand

³Biology Program, Faculty of Science Technology and Agriculture, Yala Rajabhat University, Yala, Thailand

⁴The Halal Science Center, Chulalongkorn University, Bangkok, Thailand

*Corresponding Author: imron.me@yru.ac.th, Tel.: +66-85-674-1130

Available online at: www.isroset.org

Received: 24/Aug/2022, Accepted: 21/Sept/2022, Online: 31/Oct/2022

Abstract— Purpose: The aim of this research was to preliminary study the influence of the reducing moisture method on the physicochemical properties of honey stingless bees. Research Method: By using a vacuum pump combined with a desiccator for 0, 1, 3, and 5 hours. After reducing moisture content, the honey of stingless bees was determined physicochemical properties, moisture content, pH, conductivity, Hydroxymethylfurfural (HMF) content, °Brix, total reducing sugar content, and color according to AOAC and International Honey Commission. Findings: The results showed that at 5 hours displayed the moisture content lower than others. Additionally, their physicochemical properties were nearly the raw honey stingless bee before reducing moisture content and accordingly to stingless bee honey standard. Originality/value: Research findings could be used in the agricultural or food industries for honey storage and quality control.

Keywords— Honey stingless bee; Physicochemical properties; Vacuum drying method

I. INTRODUCTION

The Stingless bee plays a significant part in plants' pollination process and an insect belongs to the order Hymenoptera of the family Apidae (Meliponini). it was found in 10 genera with 31 species in Thailand [1],[2],[3]. Honey of stingless bees is including of a diverse component of carbohydrates, such as glucose and fructose, amino acids, organic acids, vitamins, minerals, and enzymes from nature. Additionally, various bioactive and phytochemical contents for example phenolic and flavonoids [4],[5]. In the same way, it also has biological activity consisting of cardioprotective, antiinflammatory, antihelminthic, neuromodulatory action, and antioxidant activity [4]. However, honey stingless bees have a high moisture content, which results in a fermentation process and can decrease honey quality [6]. As previously reported, reducing honey's moisture by heating method might cause loss or change of components in honey [7,8]. Besides, in Thailand, the quality of honey required by the Thai Agricultural Standard of moisture content with less than 21% [9]. Therefore, finding a method that is free from heat may help with the effect of altering the honey quality.

II. RELATED WORK

In the previous research work, Kunjet et al. studied the effect of reducing moisture to maintain the honey quality of stingless bee by using the drying oven method. The data

showed that the drying of honey stingless at 40 and 60 degrees Celsius was able to maintain the hydroxymethyl furfural (HMF) and moisture content of less than 80 mg/kg and 22%, respectively [9]. However, this method was used for about 6 hrs. to reduce the moisture content to 20-21%. Moreover, the microwave vacuum drying method has been reported that obtained quality dried honey after drying and a short time process (0-2 min) at low power [10]. However, low-cost entrepreneurs or farmers may not be able to prepare with this method. Thus, we are interested in a preliminary study of the effect of the reducing moisture method on the physicochemical properties by using a vacuum pump combined with a desiccator, the basic equipment in the general laboratory.

III. METHODOLOGY

Collection and Preparation of Honey of Stingless Bee

The honey of stingless bee (*Tetragonula laeviceps*) was collected stingless bee farm in Pase Yawo sub-district, Sai Buri district, Pattani, Thailand, on February 2020. Honey was filtrated with thin whites and kept in a glass bottom at 4°C for further study.

Experimental Study

Briefly, 100 g of stingless bee honey was added into the stainless-steel chamber and taken into the desiccator, which was connected with a vacuum pump (LAX®, S LX 115 Model, 1.5 CFM, 5 Pa, 220V~50Hz, ¼ HP), and decreasing the moisture content for 0, 1, 3 and 5 hrs.

After decreasing moisture content, stingless bee honey was analyzed for its physicochemical properties. Moisture content and °Brix were conducted by using a portable refractometer. HMF content and total reducing sugar content followed the white method [11]. In brief, the pH value is determined with pH meter METTLER TOLEDO.

The concentration of the honey solution was prepared as 10% (w/v) in DI water. The conductivity value is measured with a Conductivity meter HACH Sension 378. The concentration of the honey solution was prepared as a 20% (w/v) in DI, and it was expressed in microSiemens per centimeter (S/cm) [12], [13], [14]. The color of honey was analyzed with Jasco V-730 Spectrophotometry at 560 nm. The absorbance of the sample was multiplied by a factor of 3.15. It was expressed in mm Pfund scale by compared color designations of honey in Table 1 [15].

Table 1: Color designations of honey

USDA color standard designation	Color range Pfund scale (mm)	Sample result range
Water White	≤8	0 - 0.094
Extra White	> 8 and ≤ 17	0.094 – 0.189
White	> 17 and ≤ 34	0.189 – 0.378
Extra Light Amber	> 34 and ≤ 50	0.378 – 0.595
Light Amber	> 50 and ≤ 85	0.595 – 1.389
Amber	> 85 and ≤ 114	1.389 – 3.008
Dark Amber	>114	> 3.008

Statistical Analysis

The values were the mean of three replicates and presented as mean ± standard deviation (n=3). One-way ANOVA used the analysis of the significant differences between values. The differences were considered significant when p<0.05.

IV. RESULTS AND DISCUSSION

This work evaluated the physicochemical properties at different times (1, 3, and 5 hours) for decreasing moisture content by using a vacuum pump combined with a desiccator. The influence of the method on the physicochemical properties showed that the lowest moisture content was five hrs. with 21% which takes less time than the method of Kunjet et al. [9] and also without heating. In contrast, all parameters were increased following decreasing times, except the pH value (Table 2 and 3). The data represented that the loss of moisture content increased the value of conductivity, °Brix, color, HMF content, and reducing sugar content [16] And also found that honey has a thicker and more viscous appearance with increasing time. Whereas pH values were decreased due to the relation of acidic compounds having increased concentrations [16]. In addition, the moisture content of honey with a vacuum system for 5 hours to affect the honey quality meets the standards of Thai Agricultural Standard (TAS 8003-2013).

Table 2: Physical properties of honey before and after reducing the moisture content

Time (hrs.)	pH	Moisture content (%)	Conductivity (µS/cm)	°Brix	Color
0	3.99±0.00 ^a	26.33±0.58 ^a	704.67±14.47 ^a	71.0±0.0 ^a	1.58±0.00 ^a
1	3.99±0.01 ^a	24.00±0.00 ^b	697.33±14.74 ^a	74.0±0.0 ^b	1.74±0.02 ^b
3	3.98±0.02 ^a	22.00±0.00 ^c	685.00±8.54 ^a	74.0±0.0 ^b	1.83±0.00 ^c
5	3.94±0.02 ^b	21.00±0.00 ^d	683.00±10.82 ^a	75.0±0.0 ^c	1.89±0.00 ^d

Note: ^{a-d}Means within each column followed by different letters are significantly different (p<0.05).

Table 3: Chemical properties of honey before and after reducing the moisture content

Times (hrs.)	HMF content (mg/100g)	Reducing sugar content (mg/100g)
0	28.30±1.20 ^a	50.30±2.08 ^a
1	30.30±0.60 ^b	52.00±0.02 ^a
3	32.30±0.60 ^c	58.00±0.03 ^b
5	35.30±1.50 ^d	65.00±0.02 ^c

Note: ^{a-d}Means within each column followed by different letters are significantly different (p<0.05).

V. CONCLUSION AND FUTURE SCOPE

The drying method of honey stingless bees by vacuum pump combined with a desiccator can reduce moisture content within five hrs., which honey quality was followed to Thai Agricultural Standard (TAS 8003-2013). This method showed a short time than previously reported [9] which uses heating and six hrs. Moreover, this method is also easy to set up and low cost when compared with microwave vacuum drying [10]. Currently, In Thailand, there are many areas of stingless bee cultivation. It may also be considered for its benefit in reducing humidity and helping preserve honey to maintain its quality. However, a vacuum pump combined with a desiccator reduces the moisture content of honey and may be required further studies in conditions of larger sample volumes in order for future industrial use.

ACKNOWLEDGMENT

We would like to thank the Chemistry program, Faculty of Science Technology and Agriculture, Yala Rajabhat University for facilitating and supporting this research's chemicals, instruments, and laboratory.

REFERENCES

- [1] B. Chuttong, Y. Chanbang, K. Sringarm, M. Burgett, "Physicochemical profiles of stingless bee (Apidae: Meliponini) honey from South East Asia (Thailand)," *Food Chemistry*, Vol. 192 (2016), pp.149-155, 2016.
- [2] T.A. Heard, "The of stingless bees in crop pollination," *Annual Review of Entomology*, Vol. 44, pp.183-206, 1999.
- [3] A. Klakaskorn, S. Wongsiri, S. Deowanish, O. Duangphakdee, "New Record of Stingless Bees (Meliponini: Trigona) in Thailand," *The Natural History Journal of Chulalongkorn University*, Vol. 5, Issue. 1, pp. 1-7, 2005.
- [4] R.G. de Oliveira, "Nutritional Composition and Medical Properties of Honey from Stingless Bees." *Nutrition and Food Science International Journal*, Vol. 7, Issue. 2, pp.1-2, 2018.
- [5] D.C.C. Lim, M.F., Abu Bakar, M., Majid, "Nutritional composition of stingless bee honey from different botanical

- origins,” *IOP Conference Series: Earth and Environmental Science*, **Vol. 269**, pp. 1-5, 2019.
- [6] S.K. Yap, N.L. Chin, Y.A. Yusof, K.Y. Chong, “Quality characteristics of dehydrated raw Kelulut honey,” *International Journal of Food Properties*, **Vol. 22, Issue. 1**, pp.556-571, 2019.
- [7] I. Singh, S. Singh, “Honey moisture reduction and its quality,” *Journal of Food Science and Technology*, **Vol. 55, Issue. 10**, pp.3861-3871, 2018.
- [8] Y. Eshete, T. Eshete, “A Review on the Effect of Processing Temperature and Time duration on Commercial Honey Quality,” *Madridge Journal of Food Technology*. **Vol. 4, Issue. 1**, pp.158-162, 2019.
- [9] S. Kunjet, N. Tedsree, T. Pumdara, S. Sutthiphon, “Effects of drying method on stable and quality of stingless bee honey,” *Khon Kaen Agriculture Journal*, **Vol. 45, Suppl.1**, pp.1355-1359, 2017.
- [10] I. Singh, S. Singh, “Honey moisture reduction and its quality,” *Journal of Food Science and Technology*, **Vol. 55, Issue. 10**, pp.3861-3871, 2018.
- [11] AOAC, “Official Methods of Analysis,” 17th Edition, The Association of Official Analytical Chemists, *Gaithersburg, MD, USA*, 2000.
- [12] S. Bogdanov, P. Martin, C. Lüllman, R. Borneck, M. Morlot, J. Lheritier, et al. “Harmonized methods of the European Honey Commission,” *Apidologie*, **28 (extra issue)**, pp.1-59, 1997.
- [13] J. Vankar, K. Tatu, R.D. Kamboj, “Monitoring Seasonal Dynamics of Physicochemical Water Quality of Chhari Dhandh Wetland, Gujarat (India),” *International Journal of Scientific Research in Biological Science*, **Vol.8, Issue.2**, pp.31-40, 2021.
- [14] S. Akbar, A. Nazir, S.A. Shah, “Analysis of Physico-Chemical and Microbiological Parameters of Rawal and Khanpur Dams,” *International Journal of Scientific Research in Chemical Science*, **Vol.9, Issue.2**, pp.1-7, 2022.
- [15] D. Frasco, “Analysis of Honey Color and HMF Content using a Genesys UV-Visible Spectrophotometer,” *Thermo Fisher Scientific*, **Madison, WI, USA**, 2018.
- [16] C.P. Panceri, T.M. Gomes, J.S. De Gois, D.L.G. Borges, M.T. Bordignon-Luiz, “Effect of dehydration process on mineral content, phenolic compounds and antioxidant activity of Cabernet Sauvignon and Merlot grapes,” *Food Research International*, **Vol. 54**, pp.1343-1350, 2013

AUTHORS PROFILE

Mr. I. Meechai pursued Bachelor of Science (Chemistry) from Yala Rajabhat University of Thailand in 2007, Master of Pharmaceutical Science from Prince of Songkla University of Thailand in year 2010 and Ph.D (Chemistry) from Walailak University of Thailand in year 2016.



He is currently working as Assistant Professor in the Programe of Chemistry, Faculty of Science Technology and Agriculture, Yala Rajabhat University, Thailand since 2017. He has published 6 research papers in reputed international journals including Scopus & Web of Science and 5 research papers in reputed national journals of TCI – Thai Journal Citation Index Centre. His main research work focuses on phytochemistry, natural product, structure elucidation and biological activity. He has 6 years of teaching experience and 5 years of Research Experience.

Miss R. Yusoh pursued Bachelor of Science (Chemistry) from Yala Rajabhat University of Thailand in year 2021. She has received a bronze medal in the Young Rising Stars of Science Award 2020 contest, Thailand and filed 1 petty patent in 2021.



Mr. I. Chelong pursued Bachelor of Science (Biology) from Prince of Songkla University of Thailand in 2004, Master of Science (Plant Science) from Prince of Songkla University of Thailand in year 2007 and Ph.D (Plant Science) from Prince of Songkla University of Thailand in year 2015. He is currently working as Assistant Professor in the Programe of Biology, Faculty of Science Technology and Agriculture, Yala Rajabhat University, Thailand since 2008. He has published more than 5 research papers in reputed international journals including Scopus & Web of Science and more than 5 research papers in reputed national journals of TCI – Thai Journal Citation Index Centre. His main research work focuses on Biodiversity, Climate change and phenomenology, Biological agents, and Stingless bees. He has 15 years of teaching experience and 15 years of Research Experience.



Mr. A. Sukswan. He received his B. Sc. in Chemistry (First Class Honor) in 2006 and his Ph. D. in Pharmaceutical Science in 2015 from Prince of Songkla University, Thailand. He is currently working as a researcher at The Halal Science Center, Chulalongkorn University, Thailand.



He has published more than 5 research papers in reputed international journals including Scopus & Web of Science and more than 5 research papers in reputed national journals of TCI – Thai Journal Citation Index Centre. His current research focuses on Halal forensic science and innovation, analytical chemistry, and also development of molecularly imprinted materials for food, agricultural, environmental, pharmaceutical applications. He is a technical committee of Standard and Metrology Institute of Islamic Countries (SMIIC).